

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 2 of 13

Amendments to the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

Claims 1-23 cancelled.

24. (Previously presented) A method of reducing or moderating a postprandial rise in plasma glucose in a mammal comprising administering to said mammal an amylin or an amylin agonist.

25. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-J₁-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala Ser or Hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu

J₁ is Ser, Pro or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is an amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro, and K₁ is Asn; then one or more A₁ to K₁ is a D-amino acid and Z is selected from

USSFN: 10/643,681

Amdt. Dated September 14, 2004

Page 3 of 13

the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

26. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-J₁-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Ser, Pro, Leu, Ile or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when

(a) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro and K₁ is Asn; or

(b) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Ser and K₁ is Asn;

then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

27. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵I₁-J₁-Leu-Pro-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 4 of 13

wherein

A₁ is Lys, Ala, Ser or hydrogen;B₁ is Ala, Ser or Thr;C₁ is Val, Leu or Ile;D₁ is His or Arg;E₁ is Ser or Thr;F₁ is Ser, Thr, Gln or Asn;G₁ is Asn, Gln or His;H₁ is Phe, Leu or Tyr;I₁ is Ala or Pro;J₁ is Ile, Val, Ala or Leu;K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkylloxy, aryloxy or aralkylloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn H₁ is Leu, I₁ is Pro, J₁ is Val and K₁ is Asn; then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkylloxy, aryloxy or aralkylloxy.

28. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-Pro-³⁰Thr-J₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;B₁ is Ala, Ser or Thr;C₁ is Val, Leu or Ile;D₁ is His or Arg;E₁ is Ser or Thr;F₁ is Ser, Thr, Gln or Asn;G₁ is Asn, Gln or His;

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 5 of 13

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and

provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val and J₁ is Asn; then one or more of A₁ to J₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

29. (Previously presented) The method of claim 24 wherein said amylin agonist is any one of ¹⁸Arg^{25,28}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28}Pro-h-amylin, ^{25,28,29}Pro-h-amylin, des-¹Lys^{25,28,29}Pro-h-amylin, ¹⁸Arg^{25,28,29}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28,29}Pro-h-amylin, ²⁵Pro²⁶Val^{28,29}Pro-h-amylin, or des-¹Lys²⁵Pro²⁶Val^{28,29}Pro-h-amylin.

30. (Previously presented) The method of claim 24 wherein the amylin agonist is ^{25,28,29}Pro-h-amylin.

31. (Previously presented) A method of treating ingestion of a toxin in a mammal comprising administering to said mammal an amylin or an amylin agonist and aspirating the toxin out of a stomach of the mammal.

32. (Previously presented) The method of claim 31 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-J₁-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala Ser or Hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 6 of 13

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu

J₁ is Ser, Pro or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is an amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro, and K₁ is Asn; then one or more A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

33. (Previously presented) The method of claim 31 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phc-Leu-C₁-D₁-E₁-²⁰F₁-
G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-J₁-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Ser, Pro, Leu, Ile or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 7 of 13

(c) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro and K₁ is Asn; or

(d) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Ser and K₁ is Asn;

then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

34. (Previously presented) The method of claim 31 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵I₁-J₁-Leu-Pro-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ala or Pro;

J₁ is Ile, Val, Ala or Leu;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val and K₁ is Asn; then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

35. (Previously presented) The method of claim 31 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 8 of 13

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-Pro-³⁰Thr-J₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and

provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val and J₁ is Asn; then one or more of A₁ to J₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

36. (Previously presented) The method of claim 31 wherein said amylin agonist is any one of ¹⁸Arg^{25,28}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28}Pro-h-amylin, ^{25,28,29}Pro-h-amylin, des-¹Lys^{25,28,29}Pro-h-amylin, ¹⁸Arg^{25,28,29}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28,29}Pro-h-amylin, ²⁵Pro²⁶Val^{28,29}Pro-h-amylin, or des-¹Lys²⁵Pro²⁶Val^{28,29}Pro-h-amylin.

37. (Previously presented) The method of claim 31 wherein the amylin agonist is ^{25,28,29}Pro-h-amylin.

38. (New) The method of claim 24 wherein the mammal has diabetes.

39. (New) The method of claim 38 wherein the diabetes is type 1.

40. (New) The method of claim 38 wherein the diabetes is type 2.

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 9 of 13

41. (New) The method of claim 25 wherein the mammal has diabetes.
42. (New) The method of claim 41 wherein the diabetes is type 1.
43. (New) The method of claim 41 wherein the diabetes is type 2.
44. (New) The method of claim 26 wherein the mammal has diabetes.
45. (New) The method of claim 44 wherein the diabetes is type 1.
46. (New) The method of claim 44 wherein the diabetes is type 2.
47. (New) The method of claim 27 wherein the mammal has diabetes.
48. (New) The method of claim 47 wherein the diabetes is type 1.
49. (New) The method of claim 47 wherein the diabetes is type 2.
50. (New) The method of claim 28 wherein the mammal has diabetes.
51. (New) The method of claim 50 wherein the diabetes is type 1.
52. (New) The method of claim 50 wherein the diabetes is type 2.
53. (New) The method of claim 30 wherein the mammal has diabetes.
54. (New) The method of claim 53 wherein the diabetes is type 1.
55. (New) The method of claim 53 wherein the diabetes is type 2.
56. (New) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵I₁-J₁-Leu-K₁-L₁-³⁰Thr-M₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z.

wherein

A₁ is Lys, Ala, Ser, Hydrogen or acetylated Lys;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 10 of 13

H₁ is Phe, Leu or Tyr;

I₁ is Ala or Pro;

J₁ is Ile, Val, Ala or Leu

K₁ is Ser, Pro, Leu, Ile or Thr;

L₁ is Ser, Pro or Thr;

M₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is an amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that

(a) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Phe, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Ser, and M₁ is Asn;

(b) when A₁ is Lys, B₁ is Ala, C₁ is Ile, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Pro, and M₁ is Asn;

(c) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Thr, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Pro, and M₁ is Asn;

(d) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val, K₁ is Pro, L₁ is Pro, and M₁ is Asn;

(e) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val, K₁ is Ser, L₁ is Pro and M₁ is Asn; or

(f) when A₁ is Lys, B₁ is Thr, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is His, H₁ is Leu, I₁ is Ala, J₁ is Ala, K₁ is Leu, L₁ is Pro and M₁ is Asp;

then one or more of any of A₁ to M₁ is not an L-amino acid and Z is not amino.

57. (New) The method of claim 56 wherein the mammal has diabetes.

58. (New) The method of claim 57 wherein the diabetes is type 1.

59. (New) The method of claim 57 wherein the diabetes is type 2.

60. (New) A method of reducing gastric motility or delaying gastric emptying in a mammal comprising administering to said mammal a therapeutically effective amount of an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵I₁-J₁-Leu-K₁-L₁-³⁰Thr-M₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser, Hydrogen or acetylated Lys;

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 11 of 13

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ala or Pro;

J₁ is Ile, Val, Ala or Leu

K₁ is Ser, Pro, Leu, Ile or Thr;

L₁ is Ser, Pro or Thr;

M₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is an amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that

(a) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Phe, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Ser, and M₁ is Asn;

(b) when A₁ is Lys, B₁ is Ala, C₁ is Ile, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Pro, and M₁ is Asn;

(c) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Thr, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Pro, and M₁ is Asn;

(d) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val, K₁ is Pro, L₁ is Pro, and M₁ is Asn;

(e) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val, K₁ is Ser, L₁ is Pro and M₁ is Asn; or

(f) when A₁ is Lys, B₁ is Thr, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is His, H₁ is Leu, I₁ is Ala, J₁ is Ala, K₁ is Leu, L₁ is Pro and M₁ is Asp;

then one or more of any of A₁ to M₁ is not an L-amino acid and Z is not amino.

61. (New) The method of claim 60 wherein the mammal has diabetes.

62. (New) The method of claim 61 wherein the diabetes is type 1.

63. (New) The method of claim 61 wherein the diabetes is type 2.

USSN: 10/643,681

Amdt. Dated September 14, 2004

Page 12 of 13

64. (New) The method of claim 60 wherein I₁ and K₁ are Pro and the mammal has diabetes.

65. (New) The method of claim 60 wherein I₁ and L₁ are Pro and the mammal has diabetes.

66. (New) The method of claim 60 wherein K₁ and L₁ are Pro and the mammal has diabetes.

67. (New) The method of claim 60 wherein I₁, K₁ and L₁ are Pro and the mammal has diabetes.

68. (New) The method of claim 60 wherein said amylin agonist is any one of ¹⁸Arg^{25,28}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28}Pro-h-amylin, ^{25,28,29}Pro-h-amylin, des-¹Lys^{25,28,29}Pro-h-amylin, ¹⁸Arg^{25,28,29}Pro-h-amylin, des-¹Lys¹⁸Arg^{25,28,29}Pro-h-amylin, ²⁵Pro²⁶Val^{28,29}Pro-h-amylin, or des-¹Lys²⁵Pro²⁶Val^{28,29}Pro-h-amylin and the mammal has diabetes.

69. (Previously presented) The method of claim 60 wherein the amylin agonist is ^{25,28,29}Pro-h-amylin and the mammal has diabetes.